

*Scientific Note/Comunicação Científica*

# New records of *Dictyopsocus pennicornis* (Burmeister) (Psocodea 'Psocoptera': Psocidae: Psocinae)

Registered on ZooBank: [urn:lsid:zoobank.org/pub:3DAF005A-AFC0-4D47-AA2C-2459CE397891](https://urn.isid:zoobank.org/pub:3DAF005A-AFC0-4D47-AA2C-2459CE397891)

João Alves Oliveira<sup>1</sup>, Alberto Moreira Silva-Neto<sup>1</sup>✉,  
Diego Matheus Mello Mendes<sup>1</sup> & Alfonso Neri García Aldrete<sup>2</sup>

1. Instituto Nacional de Pesquisas da Amazônia/Pós graduação em Entomologia. 2. Instituto de Biología, Universidad Nacional Autónoma de México.

## *EntomoBrasilis* 10 (2): 127-130 (2017)

**Abstract.** First records of *Dictyopsocus pennicornis* (Burmeister), in the northeastern Brazilian states of Bahia, Ceará and Paraíba, are presented, with biogeographic comments.

**Keywords:** Bark-Lice; Geographic distribution; Neotropics, Psocids; Thyrsopterini.

## Novos registros de *Dictyopsocus pennicornis* (Burmeister) (Psocodea 'Psocoptera': Psocidae: Psocinae)

**Resumo.** Os primeiros registros de *Dictyopsocus pennicornis* (Burmeister), nos estados brasileiros da Bahia, Ceará e Paraíba são apresentados, com comentários biogeográficos.

**Palavras-chave:** Distribuição geográfica; Neotropical; Piolhos de cascas de árvores; Psocídeos; Thyrsopterini.

The family Psocidae includes the subfamilies Kaindipsocinae, Amphigerontiinae and Psocinae (YOSHIZAWA *et al.* 2011), the latter including five tribes, one of which is Thyrsopterini. It is divided in two groups previously known as Cerastipsocinae and Thyrsopterinae (YOSHIZAWA & JOHNSON 2008). The tribe includes 15 genera, one of which is the monotypic *Dictyopsocus* Enderlein, represented by *Dictyopsocus pennicornis* (Burmeister), easily identified by the presence of small ramifying veinlets in the central area of the forewing. *Dictyopsocus* was raised by ENDERLEIN (1901) to include the species then known as *Thyrsopterus pennicornis* Burmeister (BURMEISTER 1839). It was redescribed by New in 1973 and later, in 1978 was transferred and reduced to a subgenus of *Thyrsopterus* by the same author, with a description of a female specimen (NEW 1973, 1978). The genus was reinstated by MOCKFORD (1992). The species is widely distributed, ranging from Venezuela to Argentina, near the border with southern Brazil (LIENHARD & SMITHERS 2002). In Brazil *D. pennicornis* was recorded for the states of Amazonas and Santa Catarina (GARCÍA ALDRETE & MOCKFORD 2009). The purpose of this note is to record the presence of *D. pennicornis* in three Brazilian northeastern states, with biogeographic comments. The specimens of *D. pennicornis* for this study were recently collected during expeditions as part of the Brazilian "PPBio Semi-árido" program, and are deposited in the Johann Becker Entomological

Collection at the Museu de Zoologia da Universidade Estadual de Feira de Santana. Four specimens were collected in the States of: a) Bahia, municipality of Santa Terezinha, Serra da Jibóia, 12°51' S; 39°28' W, 10.XI.2010, 1 male, 1 female; b) Ceará, municipality of Ubajara, Parque Nacional Serra Grande, 02°49' S, 40°54' W, 1 male (Figures 1-13), and c) Paraíba, municipality of Areia, Brejo Paraibano, Reserva Pau Ferro, 26.IX.2011, 1 female (Figures 14-26). The specimens were dissected in 80% ethanol; their parts (head, right legs and wings, and genitals) were mounted in Canada balsam. Before dissecting, whole specimens were placed in 80% ethanol under a dissecting microscope, illuminated with cold, white light, and observed at 50X to record color. The images were taken with a Leica DFC295 attached at a stereoscopic microscope M205. The four specimens were collected in forest areas. Although their presence in the northeastern region of Brazil, in the semi-arid biome Caatinga, they are only known from forests enclaves, although extensive collecting was conducted in the semi-arid regions as part of the "PPBio Semi-árido" program, which aims to document the biodiversity of this biome. The absence of *D. pennicornis* in the other areas of the semi-arid region allows us to suggest that this species does not have the ability to disperse among the forest patches through the semi-arid matrix. They must have colonized those different areas when they were connected. With exception of Caracas and Argentina, all the other forests are part

### Edited by:

William Costa Rodrigues

### Article History:

Received: 13.xii.2016

Accepted: 04.vii.2017

### ✉ Corresponding author:

Alberto Moreira Silva-Neto

✉ [bio.alberto@gmail.com](mailto:bio.alberto@gmail.com)


📍 <http://orcid.org/0000-0001-6081-4397>

### Funding agencies:


↗ Instituto Nacional de Pesquisas da Amazônia, Universidad Nacional Autónoma de México

of the Amazonian rainforest, Caatinga forest enclaves or Atlantic rainforest. They were connected at different periods of time in the Tertiary (RYLANDS *et al.* 1996) and in the Quaternary (MARKS *et al.* 2002), during forest expansion periods due to global climate changes, although there are different hypotheses regarding how they were connected (BIGARELLA & ANDRADE-LIMA 1982; CRACRAFT 1994; SILVA & CASTELETTI 2003). This study indicates that *D.*

*pennicornis* and possibly other species of Psocoptera, as an old and widespread group, is a good model for biogeography studies. A study to resolve the relationship between *D. pennicornis* populations and more sampling effort might endorse the species distribution story, which would support more evidence for one of the hypotheses about the Amazonian rainforest and Atlantic rainforest connection (MELO SANTOS *et al.* 2007).



Figures 1-13. *Dictyopsocus pennicornis* (Burmeister). Male. 1. Front view of head. 2. Forewing. 3. Hindwing. 4. Right antenna. 5. Right lacinal tip. 6. Labrum. 7. Right maxillary palp. 8. Right foreleg. 9. Right midleg. 10. Right hindleg. 11. Clunium, right paraproct, and epiproct. 12. Hypandrium. 13. Phallosome. Scales in mm.



Figures 14-26. *Dictyopsocus pennicornis* (Burmeister). Female. 14. Front view of head. 15. Forewing. 16. Hindwing. 17. Right antenna. 18. Right lacinial tip. 19. Labrum. 20. Right maxillary palp. 21. Right foreleg. 22. Right midleg. 23. Right hindleg. 24. Clunium, right paraproct and epiprocto. 25. Gonapophyses. 26. Subgenital plate.

## ACKNOWLEDGEMENTS

We thank Freddy Bravo, Coordinator of the Invertebrates Area for the PPBio-Semi-arid Project (process-558317/2009-0), at the Universidade Estadual de Feira de Santana, Bahia, Brazil. ANGA thanks Instituto de Biología, Universidad Nacional Autónoma de México, for continuous research support. AMSN, JRAO and DMMM thank Instituto Nacional de Pesquisas da Amazônia (INPA) for research support.

## REFERENCES

- Bigarella, J.J. & D. Andrade-Lima, 1982. Paleo environmental changes in Brazil, p. 27-40. *In*: Prance, G.T. (Ed). Biological Diversification in the Tropics. Columbia University Press, New York, 714 p.
- Burmeister, H., 1839. Handbuch der Entomologie. 2. Band. Berlin. XII+1050 pp. (Psocoptera: p. 772-782).
- Cracraft, J., 1994. Species diversity, biogeography, and the evolution of biotas. *American Zoologist*, 34: 33-47. DOI: <https://doi.org/10.1093/icb/34.1.33>.
- Enderlein, G., 1901. Neue deutsche und exotische Psociden, sowie Bemerkungen zur Systematik. *Zoologische Jahrbucher (Abteilung Systematik)*, 14: 537-548.
- García Aldrete, A.N. & E.L. Mockford, 2009. A list of Psocoptera (Insecta: Psocodea) from Brazil. *Revista Mexicana de Biodiversidad*, 80: 665-673.
- Lienhard, C. & C.N. Smithers, 2002. Psocoptera (Insecta). *World Catalogue and Bibliography. Instrumenta Biodiversitatis V. Muséum d'histoire naturelle, Genève, Suisse*, 745 p.
- Marks, B.D., S.J. Hackett & A.P. Capparella, 2002. Historical relationships among Neotropical lowland forest areas of endemism as determined by mitochondrial DNA sequence variation within the Wedge-billed Woodcreeper (Aves: Dendrocolaptidae: *Glyphorynchus spirurus*). *Molecular Phylogenetics and Evolution*, 24: 153-167. DOI: [https://doi.org/10.1016/S1055-7903\(02\)00233-6](https://doi.org/10.1016/S1055-7903(02)00233-6).
- Melo Santos, A.M., D.R. Cavalcanti, J.M.C.D. Silva & M. Tabarelli, (2007). Biogeographical relationships among tropical forests in north-eastern Brazil. *Journal of Biogeography*, 34: 437-446. DOI: <https://doi.org/10.1111/j.1365-2699.2006.01604.x>.
- Mockford, E.L., 1992. Taxonomy of the Thyrsophorine barklice of Panama (Psocoptera: Psocidae: Thyrsophorinae). p. 257-270. *In*: Quintero D. & Aiello A. (eds). *Insects of Panama and Mesoamerica*. Oxford University Press, Oxford, 692 p.
- New T., 1978. An appraisal of the Thyrsophorinae stat. nov. (Psocoptera, Psocidae) and of its constituent genera from the Neotropics. *Systematic Entomology* 3: 35-49. DOI: <https://doi.org/10.1111/j.1365-3113.1978.tb00386.x>.
- New, T., 1973. A note on the genus *Dictyopsocus* Enderlein (Psocoptera, Thyrsophoridae). *Journal of Natural History*, 7: 509-512. DOI: <https://doi.org/10.1080/00222937300770401>.
- Rylands A.B., G.A.B. Fonseca, Y.L.R. Leite & R.A. Mittermeier, 1996. Primates of the Atlantic Forest: origin, distributions, endemism, and communities. p. 21-51. *In*: Norconk M.A., A.L. Rosenberger, & P. A. Garber (eds). *Adaptive radiations of neotropical primates*. Plenum, New York, 348 p.
- Silva, J.M.C. & C.H.M. Casteleti, 2003. Status of the biodiversity of the Atlantic Forest of Brazil. p. 43-59. *In*: Silva, J.M.C. & C.H.M. Casteleti, (eds). *The Atlantic Forest of South America: Biodiversity Status, Threats, and Outlook*. CABS and Island Press, Washington, 488 p.
- Yoshizawa, K. & K.P. Johnson, 2008. Molecular systematics of the barklouse family Psocidae (Insecta: Psocodea: 'Psocoptera') and implications for morphological and behavioral evolution. *Molecular Phylogenetics and Evolution* 46: 547-559. DOI: <https://doi.org/10.1016/j.ympev.2007.07.011>.
- Yoshizawa, K., E. Bess & K.P. Johnson, 2011. Kaindipsocinae is a sister taxon to the rest of Psocidae (Insecta: Psocodea: 'Psocoptera'). *Invertebrate Systematics* 25: 81-90. DOI: <https://doi.org/10.1071/is11004>.

\*\*\*\*\*

### Suggestion citation:

Oliveira, J.A., A.M. Silva-Neto, D.M.M. Mendes & A.N. García Aldrete, 2017. New records of *Dictyopsocus pennicornis* (Burmeister) (Psocodea 'Psocoptera': Psocidae: Psocinae). *EntomoBrasilis*, 10 (2): 127-130.

Available on: [doi:10.12741/entomoBrasilis.v10i2.673](https://doi.org/10.12741/entomoBrasilis.v10i2.673)

