

Survey of the Hymenoptera Fauna in a “Caatinga” Area in the State of Rio Grande do Norte, Northeastern Brazil

Daniell Rodrigo Rodrigues Fernandes¹✉, Jorge Anderson Guimarães², Elton Lucio Araujo³, Rogéria Inês Rosa Lara⁴ & Nelson Wanderley Perioto⁴

1. Instituto Nacional de Pesquisas da Amazônia, e-mail: daniellrodrigo@hotmail.com (Autor para correspondência✉). 2. Embrapa Hortalícias, e-mail: jorge.anderson@embrapa.br. 3. Universidade Federal Rural do Semi-Árido, e-mail: elton@ufersa.edu.br. 4. APTA - Agência Paulista de Tecnologia dos Agronegócios Ribeirão Preto, e-mail: rirlara@yahoo.com.br, nperioto2@gmail.com.

EntomoBrasilis 7 (3): 211-215 (2014)

Abstract. The aim of this study was to carry out a survey of the fauna of the Hymenoptera families in an area of Caatinga in Mossoró, state of Rio Grande do Norte, Brazil. The samplings were carried out with Malaise traps, active between 6th of February and 6th of March, 2007 and between 13th March and 14th April, 2008. A total of 5,057 Hymenoptera, belonging to 12 superfamilies and 36 families, were collected. The most abundant superfamilies were: Chalcidoidea (1,206 specimens/23.85% of the total), Vespoidea (886/17.52%), Ichneumonoidea (837/16.55%) and Platygastroidea (801/15.84%) and the most abundant families were: Platygastridae (801/15.84%), Braconidae (616/12.18%), Pteromalidae (583/11.53%), Figitidae (454/8.98%), Pompilidae (444/8.78%) and Formicidae (268/5.30%). The data obtained allow us to affirm that the Caatinga encompasses substantial family richness of Hymenoptera and that renewed effort is necessary to sample its fauna in a more extensive way.

Keywords: Biodiversity; Faunistic inventory; Semi-arid.

Levantamento da Fauna de Hymenoptera em uma Área de Caatinga do Estado do Rio Grande do Norte, Nordeste do Brasil

Resumo. O objetivo deste estudo foi realizar levantamento das famílias de himenópteros presentes em área de Caatinga em Mossoró, Estado do Rio Grande do Norte, Brasil. As amostragens foram realizadas com armadilhas Malaise, entre 6 de fevereiro e 6 de março de 2007 e 13 de março e 14 de abril de 2008. Foram coletados 5.057 himenópteros pertencentes a 12 superfamílias e 36 famílias. As superfamílias mais abundantes foram: Chalcidoidea (1.206 exemplares/23,85% do total), Vespoidea (886/17,52%), Ichneumonoidea (837/16,55%) e Platygastroidea (801/15,84%) e as famílias mais abundantes foram: Platygastridae (801/15,84%), Braconidae (616/12,18%), Pteromalidae (583/11,53%), Figitidae (454/8,98%), Pompilidae (444/8,78%) e Formicidae (268/5,30%). Os dados obtidos demonstram grande riqueza de famílias de himenópteros na Caatinga e que estudos adicionais são necessários para estabelecer a riqueza desta fauna de forma mais extensiva.

Palavras-chave: Biodiversidade; Inventário faunístico; Semi-árido.

Hymenoptera is one of the largest and most diverse orders of insects. HANSON & GAULD (2006a) affirmed that there existed nearly 115 thousand species of Hymenoptera and estimated that such a number might reach 250 thousand; according to those authors, parasitic hymenopterans are very important for the maintenance of terrestrial biodiversity by regulating the populations of other arthropods. In the Neotropical region it is possible to find 21 superfamilies and 76 families of Hymenoptera (FERNÁNDEZ 2006); for the Brazilian territory has been reported the occurrence of 18 superfamilies and 63 families (FERNÁNDEZ & SHARKEY 2006; HANSON & GAULD 2006b).

The little that is known about Hymenoptera from Rio Grande do Norte is due to the occurrence reports in Eulophidae, Eurytomidae and Ichneumonidae (ONODY & PENTEADO-DIAS 2002; FERNANDES et al. 2012, 2014a), of braconids and eupluids associated with fruit flies (ARAUJO & ZUCCHI 2002; COSTA et al. 2005), of parasitic braconids of the leafminer fly (ARAUJO et al. 2007) and of aphids (MACEDO et al. 2010). Some authors have made checklists, such as the species of Apidae (ZANELLA 2000; SILVA 2014), Chrysidae (ZANELLA & LUCENA 2014), Ichneumonoidea (SHIMBORI et al. 2014), Ophioninae (Ichneumonidae) (FERNANDES et al. 2014b) and Vespinae (Vespidae) (ANDENA & CARPENTER 2014). PENTEADO-DIAS & SCATOLINI (2003), PENTEADO-DIAS et al. (2007) and FERNANDES et al. (2012) described new species from material collected in that state.

The Caatinga biome covers 10% of the Brazilian territory; with an area of nearly 844 thousand km² it is mainly located in the Northeast region of Brazil. It is the only biome that is exclusively Brazilian and only nearly 2% of its territory is protected as preservation areas (LEAL et al. 2005; IBGE 2014). The state of Rio Grande do Norte has nearly 95% of its area covered with Caatinga vegetation. Development of studies in the Caatinga biomes is scarce. Thus, little is known about the entomofauna associated with that type of vegetation.

The aim of this study was to carry out a survey of the fauna of Hymenoptera families present in an area of Caatinga at Mossoró, state of Rio Grande do Norte, Brazil.

MATERIAL AND METHODS

The samplings were carried out in the Santa Júlia's Farm (5°01'25"S and 37°22'57"W), located in Mossoró, state of Rio Grande do Norte, Brazil, in a Caatinga native vegetation fragment adjacent to a melon crop *Cucumis melo* L. (Cucurbitaceae). In the Caatinga fragment were set up two Malaise traps, approximately 100 m apart and 100 m of the board (melon crop). The traps were active between 6th of February and 6th March, 2007 (sample 1)

Funding Agencies: CNPq; FAPESP; CAPES.

and between 13th March and 14th April, 2008 (sample 2), both periods during the rainy season.

Hymenoptera were identified according to FERNÁNDEZ & SHARKEY (2006). For Apoidea the nomenclature follows MELO & GONÇALVES (2005), with Apidae (*sensu lato*). Posteriorly, the status of superfamilies and families follow SHARKEY (2007), except Apidae (*sensu lato*).

The studied material was deposited in the Coleção Entomológica do Laboratório de Sistemática e Bioecologia de Parasitoides e

Predadores (LRRP) of the APTA Ribeirão Preto, Ribeirão Preto, São Paulo State, Brazil (N.W. Perioto, curator).

RESULTS AND DISCUSSION

A total of 5,057 Hymenoptera were collected, belonging to 12 superfamilies and 36 families (Table 1). Six of those families do not present parasitic habits: Apidae (some are kleptoparasitic bees), Crabronidae (predators, with some species kleptoparasites), Formicidae (predators, micetophagous etc), Vespidae (predators), Argidae and Pergidae (phytophagous).

Table 1. Number and percentage of relative abundance of Hymenoptera collected with a Malaise trap in an area of Caatinga, in Mossoró, RN, Brazil, between 6th February and 6th March, 2007 (sample 1) and between 13th March and 14th April, 2008 (sample 2).

Superfamily/Family	Sample 1	Sample 2	Total	RAHt	RAHs
Apoidea	127	60	187	3.70	-
Apidae (<i>sensu lato</i>)	37	10	47	0.93	25.13
Crabronidae	90	50	140	2.77	74.87
Ceraphronoidea	3	32	35	0.69	-
Ceraphronidae	3	32	35	0.69	100.00
Chalcidoidea	385	821	1,206	23.85	-
Aphelinidae	1	13	14	0.28	1.16
Chalcididae	96	151	247	4.88	20.48
Encyrtidae	11	46	57	1.13	4.73
Eucharitidae	2	0	2	0.04	0.17
Eulophidae	43	71	114	2.25	9.45
Eupelmidae	16	16	32	0.63	2.65
Eurytomidae	26	15	41	0.81	3.40
Leucospidae	4	0	4	0.08	0.33
Mymaridae	18	53	71	1.40	5.89
Perilampidae	8	2	10	0.20	0.83
Pteromalidae	150	433	583	11.53	48.34
Signiphoridae	1	2	3	0.06	0.25
Torymidae	9	8	17	0.34	1.41
Trichogrammatidae	0	11	11	0.22	0.91
Cynipoidea	167	287	454	8.98	-
Figitidae	167	287	454	8.98	100.00
Chrysidioidea	264	237	501	9.91	-
Bethylidae	71	99	170	3.36	33.93
Chrysididae	33	37	70	1.38	13.97
Dryinidae	152	91	243	4.81	48.50
Sclerogibbidae	8	10	18	0.36	3.59
Diaprioidea	0	2	2	0.04	-
Diapriidae	0	2	2	0.04	66.67
Evanoidea	8	105	113	2.23	-
Evaniidae	8	105	113	2.23	100.00
Ichneumonoidea	361	476	837	16.55	-
Braconidae	301	315	616	12.18	73.60
Ichneumonidae	60	161	221	4.37	26.40
Platygastroidea	39	762	801	15.84	-
Platygastridae	39	762	801	15.84	100.00
Proctotruopoidea	0	1	1	0.02	-
Proctotrupidae	0	1	1	0.02	100.00
Vespoidea	597	289	886	17.52	-
Formicidae	117	151	268	5.29	30.25
Mutillidae	100	8	108	2.14	12.19
Pompilidae	367	77	444	8.78	50.11

To be continued...

Table 1. Continued...

Superfamily/Family	Sample 1	Sample 2	Total	RAHt	RAHs
Rhopalosomatidae	1	45	46	0.91	5.20
Tiphidae	1	0	1	0.02	0.11
Vespidae	11	8	19	0.38	2.14
Tenthredinoidea	23	11	34	0.67	-
Argidae	19	7	26	0.51	76.47
Pergidae	4	4	8	0.16	23.53
Total	1,974	3,083	5,057		

RAHt = percentage of relative abundance of the family or superfamily in relation to the total of Hymenoptera.

RAHs = percentage of relative abundance of the Hymenoptera families in relation to those collected in the superfamily to which it belongs.

In the Neotropical region are recorded 21 superfamilies and 72 families of Hymenoptera. The number reaches 76 if considered Andrenidae, Colletidae, Halictidae and Megachilidae (*Apidae sensu lato*) (FERNÁNDEZ 2006). For Brazil are reported 18 superfamilies and 63 families (FERNÁNDEZ & SHARKEY 2006; HANSON & GAULD 2006b). Thus, even from brief surveys of fauna, like the one here reported, it was possible to establish that 57.1% of the superfamilies and 50% of the families known for the Neotropics, 66.7% of the superfamilies and 57.1% of the families are already known for Brazil (Table 2).

Of the 5,057 Hymenoptera specimens collected in this study, 1,206 (23.85% of the total) are Chalcidoidea, a group in which Pteromalidae (583 specimens/11.53% of the total), Chalcididae (247/4.88%) and Eulophidae (114/2.25%) stand out, and which, together, represented more than 75% of the collected Chalcidoidea (Table 1). NOYES (2003) reported the existence of nearly 22,000 species of Chalcidoidea in the world. Such insects are relatively little studied. Nearly 12,000 species of insects are recorded as their hosts, which demonstrate their great capacity for parasitism and their importance as a factor of biotic mortality in the environment. GRISSELL & SCHAUFF (1997) asserted that Chalcidoidea have great diversity of biological habits: the group has, at least, 14 life strategies which include mainly solitary, gregarious, hyperparasitism and polyembryony.

A total of 886 specimens of Vespoidea (17.52%) were obtained, among which Pompilidae were the most collected (444/8.78%), followed by Formicidae (268/5.30%) and Mutillidae (108/2.14%). These three families together represent more than 90% of the collected Vespoidea (Table 1). Vespidae was poorly represented in the current survey (19/0.38%) and the reason for the small number of social wasps collected is unknown.

Ichneumonoidea was the third most collected superfamily, with 837 specimens (16.55%), among which 616 Braconidae (12.18%)

and 221 Ichneumonidae (4.37%), followed by Platygastroidea, with 801 specimens (15.84%) of Platygastridae.

The other sampled families presented relative frequencies lower than 10% (Table 1). Noteworthy is the identification of 18 specimens of Sclerogibbidae (Chrysidoidea), a group rarely found in samplings of parasitic Hymenoptera and, consequently, rare in entomological collections (AZEVEDO & SANTOS 2000). Other families considered rare in collections were also obtained: Rhopalosomatidae (46 specimens), Leucospidae (four) and Proctotrupidae (one).

The number of 30 families of parasitic Hymenoptera recorded to Caatinga in the state of Rio Grande do Norte is considerably close to that which occurs in other Brazilian biomes, like in the Atlantic Rainforest, in the state of Espírito Santo, where AZEVEDO & SANTOS (2000) and AZEVEDO *et al.* (2002, 2003) found 30, 28 and 35 families of Hymenoptera parasitoids, respectively. PERIOTO *et al.* (2003, 2005) recorded 23 and 26 parasitic wasp families in the Atlantic Rainforest in Ubatuba and Iguape, in the state of São Paulo, respectively. FEITOSA *et al.* (2007) recorded 25 parasitic Hymenoptera families in the Amazonian tropical rain forest at Manaus, in the state of Amazonas, and PERIOTO *et al.* (2008) found 27 parasitic Hymenoptera families in the Brazilian Savannah (Cerrado) in Luis Antonio, in the state of São Paulo. PÁDUA & ZAMPIEON (2012) found 20 parasitic Hymenoptera families in the Brazilian Savannah (Cerrado) in Delfinópolis, in the state of Minas Gerais.

It is important to point out that most of the surveys of Hymenoptera fauna carried out in Brazil have not taken into consideration the Vespoidea, Apoidea and the Symphyta, perhaps due to the complexity of their identification. The data obtained confirm that the Caatinga biome embodies substantial Hymenoptera biodiversity but that further studies to show its fauna in a more extensive way, are still necessary.

Table 2. Superfamilies and families of Hymenoptera recorded for the Neotropical region, for Brazil and for Rio Grande do Norte (RN) (present study).

Superfamily	Family	Brazil	RN	Superfamily	Family	Brazil	RN
Apoidea	Apidae (<i>sensu lato</i>)	X	X	Diaprioidea	Diapriidae	X	X
	Crabronidae	X	X		Monomachidae	X	
	Sphecidae	X		Evanoidea	Aulacidae	X	
Cephidoidea	Cephidae				Evaniiidae	X	X
Ceraphronoidea	Ceraphronidae	X	X		Gasteruptiidae	X	
	Megaspilidae	X		Ichneumonoidea	Braconidae	X	X
Chalcidoidea	Agaonidae	X			Ichneumonidae	X	X
	Aphelinidae	X	X		Megalyroidea	Megalyridae	X
	Chalcididae	X	X		Mymarommatidae	X	
	Encyrtidae	X	X		Orussidae	X	
	Eucharitidae	X	X		Pamphiliidae		

To be continued...

Table 2. Continued...

Superfamily	Family	Brazil	RN	Superfamily	Family	Brazil	RN
	Eulophidae	X	X	Platygastroidea	Platygastridae	X	X
	Eupelmidae	X	X	Proctotrupoidea	Heloridae	X	
	Eurytomidae	X	X		Pelecinidae	X	
	Leucospidae	X	X		Proctotrupidae	X	X
	Mymaridae	X	X	Siricoidea	Siricidae	X	
	Ormyridae	X		Stephanoidea	Stephanidae	X	
	Perilampidae	X	X	Tenthredinoidea	Argidae	X	X
	Pteromalidae	X	X		Cimbicidae	X	
	Rotoitidae				Diprionidae		
	Signiphoridae	X	X		Pergidae	X	X
	Tanaostigmatidae	X			Tenthredinidae	X	
	Tetracampidae			Trigonaloidea	Trigonalidae	X	
	Torymidae	X	X	Vespoidea	Bradynobaenidae		
	Trichogrammatidae	X	X		Formicidae	X	X
Chrysidoidea	Bethylidae	X	X		Mutillidae	X	X
	Chrysididae	X	X		Pompilidae	X	X
	Dryinidae	X	X		Rhopalosomatidae	X	X
	Embolemidae	X			Sapygidae	X	
	Plumariidae	X			Scoliidae	X	
	Sclerogibbidae	X	X		Sierolomorphidae		
	Scolebythidae	X			Tiphiidae	X	X
Cynipoidea	Cynipidae	X			Vespidae	X	X
	Figitidae	X	X	Xiphydrioidea	Xiphydriidae	X	
	Ibaliidae	X		Xyeloidea	Xyelidae		
	Liopteridae	X					

ACKNOWLEDGEMENTS

We are grateful to Alexandre Carlos Menezes Netto, Hailson Alves Ferreira, Leandro Delalibera Geremias, Leonardo de Souza Freitas, Luciano Pacelli Medeiros de Macedo and Marcos Aurélio Araujo Lima for the assistance with the collection of the material. We are also grateful to the owners of the Fazenda Santa Júlia for their permission to use the area for the collections. We thank CNPq for the undergraduate research grant for the first author. We also thank the Instituto Nacional de Ciência e Tecnologia dos Hymenoptera Parasitóides da Região Sudeste Brasileira (HYMPAR/Sudeste - CNPq/FAPESP/CAPES).

REFERENCES

- Andena, S.R. & J.M. Carpenter, 2014. Checklist das espécies de Polistinae (Hymenoptera, Vespidae) do semiárido brasileiro, p. 169-180. In: Bravo, F. & A. Calor (eds) Artrópodes do Semiárido: Biodiversidade e Conservação. Feira de Santana: Print Mídia, 298 p.
- Araujo, E.L. & R.A. Zucchi, 2002. Parasitóides (Hymenoptera: Braconidae) de moscas-das-frutas (Diptera: Tephritidae) na região de Mossoró/Assu, estado do Rio Grande do Norte. Arquivos do Instituto Biológico, 69: 65-68.
- Araujo, E.L., D.R.R. Fernandes, L.D. Geremias, A.C. Menezes-Netto & M.A. Filgueira, 2007. Mosca minadora associada à cultura do meloeiro no Semi-Árido do Rio Grande do Norte. Revista Caatinga, 20: 210-212.
- Azevedo, C.O. & H.S. Santos, 2000. Perfil da fauna de himenópteros parasitóides (Insecta, Hymenoptera) em uma área de Mata Atlântica da Reserva Biológica de Duas Bocas, Cariacica, ES, Brasil. Boletim do Museu de Biologia Mello Leitão, 11/12: 116-126.
- Azevedo, C.O., R. Kawada, M.T. Tavares & N.W. Perioto, 2002. Perfil da fauna de himenópteros parasitóides (Insecta,
- Hymenoptera) em uma área de Mata Atlântica do Parque Estadual da Fonte Grande, Vitória, ES, Brasil. Revista Brasileira de Entomologia, 46: 133-137.
- Azevedo, C.O., M.S. Corrêa, F.T. Gobbi, R. Kawada, G.O. Lanes, A.R. Moreira, E.S. Redighieri, L.M. Santos & C. Walchert, 2003. Perfil das famílias de vespas parasitóides (Hymenoptera) em uma área de Mata Atlântica da Estação Biológica de Santa Lúcia, Santa Tereza, ES, Brasil. Boletim do Museu de Biologia Mello Leitão, 16: 39-46.
- Costa, V.A., E.L. Araujo, J.A. Guimarães, A.S. Nascimento & J. LaSalle, 2005. Redescoberta de *Tetrastichus giffardianus* (Hymenoptera: Eulophidae) após 60 anos da sua introdução no Brasil. Arquivos do Instituto Biológico, 72: 539-541.
- Feitosa, M.F.B., R.B. Querino & A.L. Henriques, 2007. Perfil da fauna de vespas parasitóides (Insecta: Hymenoptera) em reserva florestal na Amazônia, Amazonas, Brasil. Entomotropica 22: 37-43.
- Fernández, F., 2006. Sistemática de los himenópteros de la Región Neotropical: estado del conocimiento y perspectivas, p. 7-35. In: Fernández, F. & M.J. Sharkey (eds) Introducción a los Hymenoptera de la Región Neotropical. Bogotá, Sociedad Colombiana de Entomología y Universidad Nacional de Colombia, 894 p.
- Fernández, F. & M.J. Sharkey, 2006. Introducción a los Hymenoptera de la Región Neotropical. Bogotá, Sociedad Colombiana de Entomología y Universidad Nacional de Colombia, 894 p.
- Fernandes, D.R.R., R.I.R. Lara & N.W. Perioto, 2012. A new species of *Symbra* (Hymenoptera, Eurytomidae, Heimbrinae) from dry forest in Brazil and new occurrence records for other Heimbrinae. Revista Brasileira de Entomologia, 56: 415-418.
- Fernandes, D.R.R., K. Schoeninger, R.I.R. Lara & N.W. Perioto, 2014a. *Henryana magnifica* Yoshimoto (Hymenoptera: Eulophidae) in Brazil: New records for the Amazon Rainforest and Brazilian Dry Forest. EntomoBrasilis, 7: 241-243.

- Fernandes, D.R.R., H.C. Onody, R.I.R. Lara & N.W. Perioto, 2014b. Annotated checklist of Brazilian Ophioninae (Hymenoptera: Ichneumonidae). EntomoBrasilis, 7: 124-133.
- Grissell, E.E. & M.E. Schauff, 1997. A handbook of the families of Nearctic Chalcidoidea (Hymenoptera). Washington, The Entomological Society of Washington, 87p.
- Hanson, P.E. & I.D. Gauld, 2006a. Introducción, p: 1-11. In: Hanson, P.E. & I.D. Gauld (eds.). Hymenoptera de la Region Neotropical Memoirs of the American Entomological Institute, 77: 994.
- Hanson, P.E. & I.D. Gauld, 2006b. Hymenoptera de la Región Neotropical. Memoirs of the American Entomological Institute, 77: 994.
- IBGE (Instituto Brasileiro de Geografia e Estatística), 2014. Mapa de biomas e de vegetação. Available in: <<http://www.ibge.gov.br/home/presidencia/noticias/21052004biomashtml.shtml>> [Accessed: 09.10.2014].
- Leal, I.R., M. Tabarelli, J.M.C. Silva, 2005. Ecologia e conservação da Caatinga: uma introdução ao desafio, p. XII-XVI. In: Leal, I.R., M. Tabarelli & J.M.C. Silva (Eds.). Ecologia e conservação da Caatinga. Recife, Editora Universitária da UFPE, 2ª Edição, 822p.
- Macedo, L.P.M., E.R. Moura Filho, A.S. Carvalho, C.E.S. Bezerra & L.C.P. Silveira, 2010. Occurrence of *Lysiphlebus testaceipes* parasiting *Aphis gossypii* in watermelon in the State of Rio Grande do Norte, Brazil. Ciência Rural, 40: 2030-2032.
- Melo, G.A.R. & R.B. Gonçalves, 2005. Higher-level bee classifications (Hymenoptera, Apoidea, Apidae sensu lato). Revista Brasileira de Zoologia, 22: 153-159.
- Noyes, J.S., 2003. About chalcidoids. Available in: <<http://www.nhm.ac.uk/research-curation/research/projects/chalcidoids/introduction.html>> [Accessed: 09.10.2010].
- Onody, H.C. & A.M. Penteado-Dias, 2002. Ocorrência dos Ophioninae (Hymenoptera: Ichneumonidae) em área de Caatinga, Rio Grande do Norte, Brasil. Arquivos do Instituto Biológico, 69: 241-242 (Suplemento).
- Pádua, D.G. & S.L.M. Zampieron, 2012. Inventário da fauna de Hymenoptera parasitóides coletados com redes de varredura em um fragmento da Serra da Babilônia, no Sudeste do Estado de Minas Gerais. Entomobrasilis, 5: 211-216.
- Penteado-Dias, A.M. & D. Scatolini, 2003. A new species of the genus *Plumarius* Philippi (Hymenoptera: Plumariidae) from Brazil. Zoologische Mededelingen, 77: 545-550.
- Penteado-Dias, A.M., M.A.B. Moreira & P.H.G. Zarbin, 2007. A new species of *Cervellus* Szépligeti (Hymenoptera, Braconidae, Braconinae) with biological notes. Revista Brasileira de Entomologia, 51: 8-11.
- Perioto, N.W., R.I.R. Lara & A. Selegatto, 2005. Himenópteros parasitóides (Insecta, Hymenoptera) da Mata Atlântica. II. Núcleo Grajaúna, Rio Verde da Estação Ecológica Juréia Itatins, Iguape, SP, Brasil. Arquivos do Instituto Biológico, 72: 81-55.
- Perioto, N.W., R.I.R. Lara, J.C.C. Santos & T.C. Silva, 2003. Himenópteros parasitóides (Insecta, Hymenoptera) da Mata Atlântica. I. Parque Estadual da Serra do Mar, Ubatuba, SP, Brasil. Arquivos do Instituto Biológico, 70: 441-445.
- Perioto, N.W., R.I.R. Lara, A.M. Vacari, L. Favoreto, N.F. Miranda, N.R. Chagas Filho & R. Pessoa, 2008. Diversidade de himenópteros parasitóides (Hymenoptera) na Estação Ecológica de Jataí, Luiz Antônio, SP, Brasil. Revista de Agricultura, 83: 125-135.
- Silva, E.M., 2014. Lista de espécies de Apidae (Hymenoptera) do semiárido com base na literatura especializada, p. 181-202. In: Bravo, F. & A. Calor (eds) Artrópodes do Semiárido: Biodiversidade e Conservação. Feira de Santana: Print Mídia, 298 p.
- Sharkey, M.J., 2007. Phylogeny and Classification of Hymenoptera. Zootaxa, 1668: 521-548.
- Shimbori, E.M., A.P.S. Loffredo, C.S. Castro, M.A. Bortoni & A.M. Penteado-Dias, 2014. Contribuição ao conhecimento da fauna de Ichneumonoidea (Hymenoptera) do Semiárido brasileiro, p. 139-152. In: Bravo, F. & A. Calor (eds) Artrópodes do Semiárido: Biodiversidade e Conservação. Feira de Santana: Print Mídia, 298 p.
- Zanella, F.C.V., 2000. The bees of the Caatinga (Hymenoptera, Apoidea, Apiformes): a species list and comparative notes regarding their distribution. Apidologie, 31: 579-592.
- Zanella, F.C.V. & D.A.A. Lucena, 2014. Chrysidae (Hymenoptera) do Semiárido do Nordeste do Brasil: conhecimento atual e desafios, p. 153-167. In: Bravo, F. & A. Calor (eds) Artrópodes do Semiárido: Biodiversidade e Conservação. Feira de Santana: Print Mídia, 298 p.

Suggested citation:

Fernandes, D.R.R., J.A. Guimarães, E.L. Araújo, R.I.R. Lara & N.W. Perioto, 2014. Survey of the Hymenoptera Fauna in a "Caatinga" Area in the State of Rio Grande do Norte, Northeastern Brazil. EntomoBrasilis, 7 (3): 211-215.

Available on: doi:10.12741/ebrasili.v7i3.453

